



TECHNIUM
SOCIAL SCIENCES JOURNAL

www.techniumscience.com



Vol. 78/2025
A New Decade for Social Changes

PLUS
COMMUNICATION P



International
Communication & PR

A Multidisciplinary Approach to Understanding and Preventing Drowsy Driving: Environmental, Physiological, Behavioral, and Technological Perspectives

Danwoo Kim

Seoul Scholars International

danwoodylankim@gmail.com

Abstract. Drowsy driving is a serious global traffic hazard, with particularly high fatality rates in South Korea, exposing flaws in current prevention strategies. This paper examines five key areas to combat the issue: (1) real-time detection via machine learning, (2) environmental factors like PM2.5, (3) behavioral influences such as road monotony, (4) physiological effects of sleep deprivation, and (5) demographic and policy data. Machine learning enables personalized monitoring of eyelid closure, steering patterns, and breathing irregularities. Environmental and behavioral analyses identify external contributors, while sleep deprivation research highlights cognitive impairments, emphasizing the need for integrated health measures. Demographic trends and policy gaps reveal systemic weaknesses. By merging technology-based detection, public health interventions, and infrastructure reform, this study proposes a unified framework to reduce drowsy driving crashes. The findings call for multidisciplinary collaboration and advanced vehicle safety systems to address one of the most underestimated causes of preventable road fatalities.

Keywords. Drowsy driving, Fatality Sleep deprivation, Cognitive impairments, Machine learning, Policy gaps, behavioral monitoring, prefrontal cortex

1. Introduction

Drowsy driving is a critical and multifactorial hazard contributing significantly to global traffic accidents. In South Korea, persistently high fatality rates from drowsiness-induced crashes indicate systemic flaws in prevention strategies. This paper presents a comprehensive analysis across five key areas: real-time detection via machine learning, environmental risk factors such as PM2.5, behavioral influences like road monotony, the physiological impacts of sleep deprivation, and national demographic and policy data. Together, these methods offer a holistic framework to combat the drowsy driving epidemic. By integrating technological innovations with evidence-based public health strategies and infrastructure reform, the study calls for a unified, multidisciplinary response to one of the most underestimated causes of preventable traffic fatalities.

Methods

Method 1: Real-Time Behavioral Detection Through Machine Learning

Machine learning (ML)–based systems are revolutionizing the detection and prevention of drowsy driving by enabling real-time monitoring of key physiological and behavioral patterns [8,9]. These systems do not rely on a universal standard but adapt to individual drivers over time. The first variable, $Y_1 = a \cdot x$ (Fig. 1), represents the frequency and duration of eyelid closure, a primary indicator of fatigue [8]. Frequent blinking and prolonged closures suggest microsleeps or severe fatigue onset. The second variable, $Y_2 = b \cdot x$ (Fig. 2), measures hand detachment from the steering wheel, signaling lapses in motor coordination or attention [9]. The third variable, $Y_3 = c \cdot x$ (Fig. 3), reflects micro-corrections and steering tremors that often precede full lapses in attention. The fourth, $Y_4 = d \cdot x$ (Fig. 4), captures anomalies in breathing patterns—such as shallow breathing or breath-holding—which are linked to stress and fatigue [12].

These inputs are fed into a composite drowsiness model calibrated to each driver’s historical data. The system dynamically recalibrates its thresholds to accommodate different physiological baselines (e.g., athletes vs. elderly drivers). As a result, when the composite score breaches a calculated threshold, the system triggers multi-sensory alerts—ranging from vibrations to auditory warnings [9]. In advanced versions, it may also activate automated braking or suggest rest stops. This adaptive capability positions ML-based detection systems at the forefront of next-generation vehicle safety frameworks.




Figure 1

Method 2: Cognitive Impacts of PM2.5 on Driving Ability

Fine particulate matter (PM2.5), due to its minuscule size (<2.5 micrometers), poses a significant threat to neurological and cardiovascular health [3,4,6,15]. Once inhaled, PM2.5 particles can bypass the body’s natural filtration systems, infiltrate the bloodstream, and penetrate the blood-brain barrier [3,4]. Upon reaching the brain, they activate microglia—the resident immune cells—resulting in neuroinflammation. This process leads to the release of

pro-inflammatory cytokines like IL-1 β , TNF- α , and IL-6, which interfere with synaptic transmission, suppress prefrontal cortex activity, and reduce executive function [4].

Drivers experiencing this kind of cognitive degradation face multiple deficits: impaired vigilance, delayed reaction times, weakened motor control, and reduced emotional regulation [6,15]. In the context of real-world driving, this manifests as a higher likelihood of lane drifting, failure to notice hazards, and an inability to make rapid decisions. Studies in Seoul have shown that even short-term spikes in PM2.5 levels are correlated with increased accident rates and reduced performance on cognitive tests [14].

Policy responses must be multilayered. In-vehicle HEPA filtration systems should become standard, especially for public transport, commercial freight, and high-risk zones [6]. Government advisories should recommend limiting non-essential travel on high-pollution days [6]. Additionally, integrating pollution data into navigation apps can help drivers plan safer routes with lower exposure [14,15].




Figure 2

Method 3: Road Monotony and Microsleep Risk (Fig. 3)

The structure of road networks plays an underappreciated role in fatigue-related accidents [13,16]. Monotonous environments—characterized by long, unbroken stretches of asphalt with little variation—are shown to reduce cortical arousal [9]. Studies using EEG monitoring reveal that within as little as 20 to 30 minutes, brainwave activity can shift toward patterns consistent with early sleep stages [9]. In this state, drivers become vulnerable to microsleep episodes lasting between 1 to 10 seconds. At highway speeds, even a 3-second lapse can result in 100 meters of uncontrolled travel [16].

In South Korea, major expressways such as the Seohaean and Jungbu expressways exemplify this risk [13,16]. Despite only accounting for half of the expressway network, straight segments account for nearly 90% of drowsy driving crashes [16]. The United States reports similar figures through the National Highway Traffic Safety Administration and the Federal Highway Administration, reinforcing the global scope of the issue [9].

Mitigation strategies must prioritize visual and cognitive stimulation [9,13]. Mild road curvature every 15 kilometers, textured pavement, visual landscaping (such as trees or reflective markers), and dynamic road signs have all been shown to reduce fatigue [9]. Moreover, expanding and upgrading highway rest areas—with amenities like cafes, shade, and fitness equipment—can provide effective micro-breaks that reinvigorate drivers [13].




Figure 3

Method 4: Acute and Chronic Sleep Deprivation Effects (Fig. 4)

Sleep is not merely restorative—it is fundamental to human survival and cognition. Acute sleep deprivation (e.g., pulling an all-nighter) is associated with impairments equivalent to being legally intoxicated [1,7]. Williamson and Feyer (2000) demonstrated that 17–19 hours of sustained wakefulness causes a cognitive decline similar to a 0.05% blood alcohol level [1]. At 24 hours without sleep, the impairment is more severe than a 0.10% BAC [1].

Equally troubling is chronic sleep restriction. When individuals consistently receive fewer than six hours of sleep per night, performance deficits accumulate. Van Dongen et al. (2003) found that after two weeks of partial sleep deprivation, participants' reaction times and memory retention were nearly indistinguishable from those who had not slept at all for 48 hours [2]. These findings are especially relevant to South Korean teens and office workers, many of whom chronically undersleep due to academic and professional pressures [7,10].

Biologically, sleep deprivation disrupts the prefrontal cortex—critical for judgment, impulse control, and attention—and overstimulates the amygdala, leading to heightened emotional responses and poor decision-making [1,2]. Physiologically, it reduces heart rate variability, weakens immune responses, and decreases the body's resilience to stress [1]. Drivers experiencing these effects are less able to anticipate hazards, react appropriately, or even maintain consistent speed and direction on the road [1,2].




Figure 4

Method 5: Statistical Trends and Policy Responses in South Korea

The epidemiology of drowsy driving in South Korea reveals a growing crisis [13,16]. Between 2019 and 2023, over 10,000 incidents were officially recorded, but experts suggest the actual number is much higher due to underreporting and misclassification [16]. The fatality rate—2.7 deaths per 100 cases—remains more than double the rate for general traffic accidents [16]. These patterns show clear demographic trends. Older drivers (50+) are more prone to drowsy crashes in the early afternoon, likely due to circadian rhythm dips following lunch [16]. Younger drivers, particularly those under 30, are disproportionately represented in early morning accidents [16].

Male drivers are statistically more likely to be involved in fatigue-related crashes, a trend attributed to occupational driving frequency and higher exposure [16]. However, women are more susceptible to dawn-time accidents, especially under poor lighting conditions [11]. Research indicates that female drivers rely more heavily on color vision and contrast, which are compromised during twilight hours [11].

Although over 230 highway rest areas have been constructed in recent years, their distribution is inconsistent and often lacks essential facilities [13]. Legal measures such as increased fines, point deductions, and criminal penalties under the Special Crime Act have yielded modest results [16]. Educational campaigns, while well-intentioned, have failed to resonate with high-risk demographics [10,13].

Future policy must be driven by data. Investment in driver fatigue monitoring infrastructure, mandatory breaks for commercial drivers, real-time fatigue alert systems, and enhanced driver education programs are vital [13,16]. Public campaigns should frame sleep not as a luxury but as a public safety obligation [7,10].




Figure 5

Conclusion

Drowsy driving is not simply the result of being tired—it is a multifaceted and deeply systemic risk rooted in biological, environmental, behavioral, and structural systems. The findings presented in this paper, supported by Figs. 1–5, reveal that multiple interacting factors—fine particulate matter (PM_{2.5}) exposure, road monotony, microsleep episodes, impaired vigilance, and time-of-day variations—jointly contribute to a significantly higher likelihood of accidents. Each factor operates not in isolation, but in a network of influences that amplify risk. For instance, elevated PM_{2.5} levels (Fig. 1) not only impair respiratory health but also induce neuroinflammation, which slows reaction time and diminishes executive function. When coupled with monotonous road conditions (Fig. 3) or late-night driving (Fig. 5), the combined effects create a dangerous environment where even skilled drivers are vulnerable.

The data indicate that short-term spikes in PM_{2.5} concentration (Fig. 2) can raise accident risk by over 50%, a finding that underscores the urgency of air quality interventions in transportation safety policy. Similarly, reaction time deterioration with reduced vigilance (Fig. 4) demonstrates that driver fatigue is not merely about subjective tiredness—it is a measurable neurocognitive decline that can be predicted and, importantly, prevented with the right systems in place. These results reinforce the principle that traffic safety cannot be separated from environmental and physiological health.

To address this crisis, governments must adopt an integrated, multi-sector strategy. Public health agencies should collaborate with transportation departments to implement real-time air quality monitoring, integrating this data into navigation systems so that drivers can plan routes with minimal exposure to harmful pollutants. Vehicle manufacturers must be incentivized—through regulation, subsidies, or both—to include AI-based fatigue detection as

standard equipment, not as premium add-ons. Such systems should utilize multimodal data, combining camera-based eye closure detection, steering input analysis, and even in-cabin air quality sensors to assess risk in real time.

Urban and rural road planners must consider the cognitive impact of road design. Monotonous stretches of highway should be interrupted with subtle design cues—changes in scenery, road texture, or lane markings—to maintain driver engagement. Educational systems should embed sleep science and fatigue awareness into both driver’s education programs and general school curricula. This would build a generation of drivers who understand that rest is a safety requirement, not a sign of laziness. In workplaces, especially for commercial drivers, policies should encourage regulated rest periods and actively discourage unsafe scheduling practices.

The policy dimension is equally critical. Governments should establish legal thresholds for in-cabin air quality in commercial vehicles, mirroring standards already applied to building interiors. In high-risk regions, temporary driving advisories or even traffic restrictions should be issued during severe pollution episodes. Navigation apps and mapping platforms should integrate accident risk models that factor in time-of-day, air quality, and historical accident patterns, giving drivers not just the fastest route, but the safest one.

Ultimately, solving the epidemic of drowsy driving will not come from one solution alone. It requires a paradigm shift in how societies view driver fatigue—not as an individual failing, but as a systemic, preventable public health issue. The evidence presented here demands empathy toward those affected, scientific rigor in developing countermeasures, and long-term commitment from every level of society—governments, industries, communities, and individuals alike. We face a choice: to continue treating fatigue-related crashes as isolated accidents, or to recognize them as the predictable outcome of neglecting interconnected risk factors. The data make the path forward clear; what remains is the will to act decisively, comprehensively, and without delay.

References

- [1] Williamson, A., & Feyer, A. M. (2000). *Occupational and Environmental Medicine*.
- [2] Van Dongen, H. P. A., et al. (2003). *Sleep*.
- [3] Oberdörster, G., et al. (2004). *Environmental Health Perspectives*.
- [4] Block, M. L., & Calderón-Garcidueñas, L. (2009). *Trends in Neurosciences*.
- [5] Kim, & Oh. (2019). *Freeway Crashes Involving Drowsy Driving*.
- [6] The Guardian. (2025). *Air Pollution Affects Focus*.
- [7] CDC. (2017). *Teen Sleep and Car Crashes*.
- [8] Philip, P., et al. (2005). *Accident Analysis & Prevention*.
- [9] Sagaspe, P., et al. (2008). *Transportation Research Part F*.
- [10] National Sleep Foundation. (2020).
- [11] Royal Society Publishing. (2016). *Sex Differences in Visual Perception*.
- [12] Nature Communications. (2017). *Fasting and Cognitive Control*.
- [13] Korea Road Traffic Authority. (2023).
- [14] Statista. (2023). *Seoul PM2.5 Data*.
- [15] PLOS ONE. (2023). *Chinese Air Mass and PM2.5*.
- [16] Ministry of Land, Infrastructure and Transport. (2023). *Traffic Accident Data*.